

Paleoseismology and paleotsunamis of the NE Japan subduction zone and relationships with the 2011, M9 Tohoku earthquake: Constraints on the seismic cycle (PALET)

France-Japan (ANR-JST) research cooperative 'PALET' project

France



Mustapha Meghraoui (University of Strasbourg)
Matthieu Ferry (Montpellier University)
Silke Schmidt (University of Strasbourg)
Esra Cetin (Istanbul Technical University)

Japan

Shinji Toda (IRIDeS, Tohoku University)
Hiroyuki Tsutsumi (Kyoto University)
Tsuyoshi Haraguchi (Osaka City University)
Koji Okumura (Hiroshima University)

2011 Tohoku rupture may be a “super cycle” earthquake

11 March 2011 M=9.0 Tohoku, Japan



To reveal the seismic cycle of M9 events

1. **Paleo-tsunami** record in Holocene deposits
2. Long-term behavior of directly **triggered inland normal faults**
 - Long-term deformation using **marine terraces** and comparison with the 2011 coseismic movements



Back marsh inundated by the 2011 tsunami may preserve older events



震災後の広田町大野・田谷海岸 2011/06/09 9:41:32

「東日本大震災大船渡市・陸前高田市記録写真集 気仙の惨状」村田プリントサービス (2011) による


Back marsh inundated by the 2011 tsunami may preserve older events



Back marsh inundated by the 2011 tsunami may preserve older events




800-m-apart two cores share the same Tsunami event horizons in Rikuzen-Takata



Event MH2





800-m-apart two cores share the same Tsunami event horizons in Rikuzen-Takata



Event TY2


5 cm

Offshore tsunami deposits from Otsuchi reveal major tsunamis, possibly every hundreds of years, rather than every 1,000 yr




Haraguchi et al. (2006), Haraguchi & Ishibe (2009),
Haraguchi et al. (2012, AGU poster)


Offshore tsunami deposits from Otsuchi reveal major tsunamis, possibly every hundreds of years, rather than every 1,000 yr





Haraguchi et al. (2006), Haraguchi & Ishibe (2009),
Haraguchi et al. (2012, AGU poster)




Onshore evidence for paleo-tsunamis along Sanriku coast suggests about 500-yr recurrence of huge tsunamis








Together with the Sendai and Fukushima data, large tsunamis might have occurred every 350-550 years




Another way to infer seismic cycle of M~9 megathrusts is from paleoseismic data on triggered coastal normal faults





Another way to infer seismic cycle of M~9 megathrusts is from paleoseismic data on triggered coastal normal faults



April 11, 2012 earthquake (Mw 6.6): Normal faulting surface rupture





April 11, 2012 earthquake (Mw 6.6): Normal faulting surface rupture





Trenches across the Itozawa fault exposed evidence for the past events



Tsunagi



Trench walls at Tsunagi, northern Itozawa fault



Trench walls at Tsunagi, northern Itozawa fault



Our trench wall exposed evidence for the penultimate surface rupturing shock

East

West

1.1 m (2011 event)


Only 2011 event


post-penultimate

2011 event + more



Liquefaction

MRE: 12,600~17,400 cal.yBP







A group of normal faults
in Fukushima could
reveal M9 super cycles



Contradiction between 2011 coseismic subsidence and long-term coastal uplift



Ozawa et al., 2011



Older marine terraces (680-850 ka) are
at altitudes of 150-250 m on the northern
Sanriku coast: *Longterm uplift*



E

W


mT15, ~570 ka
161 m asl

mT17, ~690 ka
196 m asl


mT19, ~780 ka
235 m asl

mT19


Long-term uplift rate




Height and ages after Koike & Machida (2001), own uplift rate determination



Co- and postseismic vertical deformation




Postseismic after Ozawa et al. (2011) Nature





Long-term vertical deformation

- Marine benchmarks most useful
- Their formation requires a long period of stable relative sea level (e.g. sea level stability and tectonic quiescence)



After Scheffers et al. (2012) Holocene

Wave-cut notches




Wave-cut platforms


Height values are given as pre-2011 level (corrected for tide and co- and postseismic deformation)



Distribution and height of lower wave-cut platforms and terraces





Several benchmarks above sea level => Interseismic uplift



Coseismic & Interseismic deformation

Area of highest coseismic subsidence coincides with area of highest interseismic uplift



E

W


mT15, ~570 ka
161 m asl

mT17, ~690 ka
196 m asl

mT19, ~780 ka
235 m asl

mT19

Long-term uplift rate




- Pleistocene uplift rates are significantly lower than Holocene uplift rates.
- Tendency that MIS5e has lowest rates (slip rate variability with time).
- 3 areas:
 - north: 0.2-0.4 mm/a,
 - south: 0.1-0.2 mm/a,
 - fault influenced: high and variable uplift


High coseismic subsidence in same area as lower long-term rates. Different rates due to the seismic cycle?

Correlation of datasets

Seismicity, Coupling & Coseismic deformation




Millennial uplift rates




Correlation of datasets

Seismicity, Coupling & Coseismic deformation



Millennial uplift rates




Segmentation of the Japan trench:


=> Repetition of the 2011 Tohoku earthquake was & is likely.

Late Quaternary benchmarks indicate strong uplift:


=> significant interseismic deformation.



Centennial geodetic data shows continuous (interseismic?) subsidence



Nishimura (2012)



To explain the Sanriku coastal data, either:

- 1) Gradual but episodic uplift occurs between earthquakes
- 2) 2011 M9 event typical, with long-term subsidence
- 3) A hidden huge event is still needed to uplift the coast

Major Outcomes

- 350-550 years of recurrence interval of extreme large tsunamis hit the Sanriku coast
- Presumed active faults in Fukushima Hamadori region, triggered by the 2011 earthquake, are confirmed active and recurrence interval of surface rupturing earthquakes on each fault is in the order of 10,000 years.
- Marine terraces along the Tohoku pacific coast have been uplifted at 0.2-0.4 mm/yr and 0.1-0.2 mm/yr in Aomori-northern Iwate, and Iwate-Fukushima respectively during late Quaternary period. The latter region corresponds to the 2011 earthquake.
- Together with these findings, we model crustal deformation and estimate the maximum 550-year recurrence intervals of M9 class earthquakes.